arXiv:1003.4035v1 [math.SG] 21 Mar 2010

Multiple Rotation Type Solutions for Hamiltonian Systems on $T^\ell \times \mathbb{R}^{2n-\ell}$

Hui Qiao

Chen Institute of Mathematics and LPMC

Nankai University, Tianjin 300071

The People's Republic of China

E-mail: qiaohuimath@mail.nankai.edu.cn

Abstract

This paper deals with multiplicity of rotation type solutions for Hamiltonian systems on $T^\ell \times \mathbb{R}^{2n-\ell}$*. It is proved that, for every spatially periodic Hamiltonian system, i.e., the case* $\ell = n$*, there exist at least* $n+1$ *geometrically distinct rotation type solutions with given energy and rotation vector. It is also proved that, for a class of Hamiltonian systems on* $T^\ell \times \mathbb{R}^{2n-\ell}$ *with* $1 \leqslant \ell \leqslant 2n-1$ *but* $\ell \neq n$*, there exists at least one periodic solution or* $n+1$ *rotation type solutions on every contact energy hypersurface.*

2010 MSC: 37J45

Key words: Hamiltonian system; rotation type solution; contact type; symplectic dilation; periodic extension

1 Introduction and main result

Existence and multiplicity of periodic solutions of spatially periodic Hamiltonian systems had been studied in [1, 3, 4, 5, 7, 10, 11] and some references therein. This paper is motivated by [6].

We make the following assumptions about Hamiltonian function H

(H0) $H \in C^2(\mathbb{R}^{2n}, \mathbb{R})$.

(H1) $H(p, q+m) = H(p,q),\ \forall (p,q) \in \mathbb{R}^{2n-\ell} \times \mathbb{R}^{\ell},\ m \in \mathbb{Z}^{\ell},\ 1 \leqslant \ell \leqslant 2n$.

(H2) $\exists\, \mu, r > 0$ such that

$$0 < \mu H(p,q) \leqslant p \cdot H_p(p,q),\ \forall (p,q) \in \mathbb{R}^{2n-\ell} \times \mathbb{R}^{\ell},\ |p| \geqslant r.$$

Spatially periodic Hamiltonian functions are the special case of $\ell = n$ in (H1)-(H2). If $\ell = 2n$, then H only satisfies (H0) and (H1) and can be considered as a Hamiltonian function on the torus T^{2n}, on which Hamiltonian systems had been studied in [2, 3, 4]. If $1 \leqslant \ell \leqslant 2n-1$, an alternative result had been obtained by M. Y. Jiang in [9].

We look for solutions of

$$\begin{cases} \dot{z}(t) = JH'(z(t)), \\ z(T) = z(0) + (0,k), \\ H(z(t)) \equiv M, \end{cases} \tag{1.1}$$

with prescribed energy M and $k \in \mathbb{Z}^{\ell}$. A pair (z, T) satisfying (1.1) with $k \neq 0$ is called a ***rotation type solution*** with ***rotation vector*** k. Two solutions (z_1, T_1) and (z_2, T_2) of (1.1) are called ***geometrically distinct*** if

$$\pi \circ z_1(\mathbb{R}) \neq \pi \circ z_2(\mathbb{R}), \tag{1.2}$$

where π denotes the covering projection $\mathbb{R}^{2n-\ell} \times \mathbb{R}^{\ell} \to \mathbb{R}^{2n-\ell} \times \mathbb{R}^{\ell}/\mathbb{Z}^{\ell}$. Denote by $\tilde{\mathcal{P}}(M,k)$ the set of geometrically distinct solutions of (1.1). For the constant $r > 0$ in (H2), set

$$M^* := \max\{H(p,q) \mid |p| \leqslant r, q \in \mathbb{R}^{\ell}\}. \tag{1.3}$$

The main result in this paper is

Theorem 1.1. *Assume that* $H : \mathbb{R}^{2n} \to \mathbb{R}$ *satisfies (H0)-(H2) with* $\ell = n$. *For every* $M > M^*$ *and* $k \in \mathbb{Z}^n \setminus \{0\}$, *we have* ${}^{\#}\tilde{\mathcal{P}}(M,k) \geqslant n+1$.

As an application of Theorem 1.2 of [9], we have

Theorem 1.2. *Assume that* $H : \mathbb{R}^{2n} \to \mathbb{R}$ *satisfies (H0)-(H2) with* $1 \leqslant \ell \leqslant 2n-1$ *but* $\ell \neq n$*. For* $M > M^*$*, if* $H^{-1}(M)$ *is of contact type then* ${}^{\#}\tilde{\mathcal{P}}(M,0) \geqslant 1$ *or* ${}^{\#}\tilde{\mathcal{P}}(M,k_0) \geqslant n+1$ *for some* $k_0 \in \mathbb{Z}^\ell \setminus \{0\}$*.*

Remark 1.3. Assumptions of Theorem 1.1 are the same with that of Theorem 0.1 of [6] which states that ${}^{\#}\tilde{\mathcal{P}}(M,k) \geqslant n$, while Theorem 1.1 gives one more solution of (1.1). When $n = 1$, Theorem 0.1 of [6] gives one rotation type solution, while Theorem 1.1 gives two. For example, consider the simple pendulum equation $\ddot{q} + \sin(q) = 0$ with Hamiltonian function $H(p,q) = \frac{1}{2}p^2 - \cos(q)$. For every $M > 1$, the energy surface $H^{-1}(M)$ consists of two disjoint curves z_1 and z_2, each one is a rotation type solution. See Fig 1.

The idea of the proof of Theorem 1.1 is as follows. In [6], P. Felmer modified H quadratically to obtain a new Hamiltonian function $\bar{H}$ which is regular near $\bar{H}^{-1}(M) = H^{-1}(M)$, and then obtained solutions $(\bar{x}, \bar{T})$ of

$$\begin{cases} \dot{x}(t) + (0,k) = \bar{T}J\bar{H}'(x(t) + t(0,k)), \\ x(1) = x(0), \\ \bar{H}(x(t) + t(0,k)) \equiv M. \end{cases} \tag{1.4}$$

A reparametrization of $\bar{x}(t) + t(0,k)$ gives a solution of (1.1). This proves the existence. **For the multiplicity proof, instead of Ljusternik-Schnirelmann category argument in [6], we adopt a different method:** Recall that Rabinowitz' idea to construct a Hamiltonian function H_Σ for a starshaped hypersurface Σ with respect to the origin is to set

$$H_\Sigma(x) = 1, \ \forall x \in \Sigma,$$
$$H_\Sigma(\lambda x) = \varphi(\lambda), \ \forall x \in \Sigma, \ \lambda \geqslant 0,$$

where φ is some well chosen function. In this paper, we use a modification of this idea to define a Hamiltonian function $\hat{H}$ which is periodic in all components and possesses $H^{-1}(M)$ as a component of one of its regular energy hypersurfaces. From a solution of (1.4), we obtain a solution $(\hat{x}, \hat{T})$ of

$$\begin{cases} \dot{x}(t) + (0,k) = \hat{T}J\hat{H}'(x(t) + t(0,k)), \\ x(1) = x(0). \end{cases} \tag{1.5}$$

For the number $\hat{T}$, following the idea of [4], we obtain $2n+1$ solutions of (1.5). **These $2n+1$ solutions are classified into two categories and give at least $n+1$ geometrically distinct solutions of (1.1).**

To prove Theorem 1.2, define a Hamiltonian function $\breve{H}$ which possesses $H^{-1}(M)$ as a regular energy hypersurface, is periodic in q and equals to its maximum outside an open neighborhood of $H^{-1}(M)$ in $\mathbb{R}^{2n}$. By Theorem 1.2 of [9], we obtain a solution of (1.5) with k equals to some $k_0 \in \mathbb{Z}^\ell$. If $k_0 \neq 0$, extending $\breve{H}$ periodically in p and repeating the multiplicity proof of Theorem 1.1, we obtain $n+1$ geometrically distinct solutions of (1.1).

This paper is divided into four sections. In Section 2, we prepare some preliminaries and prove the existence of (1.5). In Section 3, we follow the idea of [4] to obtain the multiplicity of (1.5). Here, special care in the proof of the Corollary of Theorem 5 of [4] has to be taken. The existence proof of (1.5) in Section 2.3 yields the existence of rest points. Then, assuming the number of them is finite, they form a Morse decomposition. Hence, the existence proof of (1.5) is necessary. Finally in Section 4, we prove Theorem 1.1 and Theorem 1.2.

In this paper, we denote by $x \cdot y$ the standard inner product of x and y in Euclidean space, and S^d the unit sphere in $\mathbb{R}^{d+1}$.

2 Preliminaries

2.1 Some properties of Hamiltonian functions H satisfying (H0)-(H2)

By (H0)-(H2), the energy hypersurface $H^{-1}(M)$ is bounded in p and periodic in q.

Proposition 2.1. (cf. Lemma 1.1 of [6]) *For every $M > M^*$, the following holds.*

(i) There exists two constants $r' = r'(H,M), r'' = r''(H,M) > r$ such that

$$r' \leqslant |p| \leqslant r'', \ \forall (p,q) \in H^{-1}(M). \tag{2.1}$$

(ii) Given $p \in \mathbb{R}^{2n-\ell} \setminus \{0\}$, there exists a unique periodic function $\sigma = \sigma_p \in C^2(\mathbb{R}^\ell, \mathbb{R}_+)$ such that

$$H(\sigma(q)p, q) = M, \ \sigma(q+m) = \sigma(q), \ \forall q \in \mathbb{R}^\ell, \ m \in \mathbb{Z}^\ell. \tag{2.2}$$

(iii) The energy hypersurface $H^{-1}(M)$ is periodic in q in the sense of

$$H^{-1}(M) = H^{-1}(M) + \{0\} \times \mathbb{Z}^{\ell}. \tag{2.3}$$

Proof. Conclusions of this proposition are the same with that of Lemma 1.1 of [6] except the upperbound in (2.1) whose proof is given below. By (H2), we have

$$H(p,q) \geqslant a|p|^{\mu},\ \forall (p,q) \in \mathbb{R}^{2n-\ell} \times \mathbb{R}^{\ell},\ |p| \geqslant r, \tag{2.4}$$

where

$$a := \min_{|p|=r} \frac{H(p,q)}{|p|^{\mu}} > 0. \tag{2.5}$$

Set

$$r' := \min\{|p| \mid (p,q) \in H^{-1}(M)\},\ r'' := \left(\frac{M}{a}\right)^{\frac{1}{\mu}}. \tag{2.6}$$

By (1.3) and (2.4)-(2.6), we obtain (2.1). ■

Remark 2.2. In (H2) of [6], the condition $\mu > 1$ is assumed. However, in the proof of Lemma 1.1 of [6], condition $\mu > 0$ is sufficient for making $f'(\sigma) > 0$ and $f(\sigma) \to +\infty$ as $\sigma \to +\infty$ in (1.5) of [6].

By (1.9) of [6], there exists a unique C^2 function

$$\alpha : \mathbb{R}^{2n}_{*} := \mathbb{R}^{2n} \setminus (\{0\} \times \mathbb{R}^{\ell}) \to \mathbb{R}_{+}, \tag{2.7}$$

which is positive homogeneous of degree one in p such that

$$H\left(\frac{p}{\alpha(p,q)}, q\right) \equiv M \quad \text{and} \quad \alpha(p, q+m) = \alpha(p,q),\ \forall (p,q) \in \mathbb{R}^{2n}_{*},\ m \in \mathbb{Z}^{\ell}. \tag{2.8}$$

In fact, for any $\lambda > 0$, we have

$$M = H\left(\frac{\lambda p}{\alpha(\lambda p, q)}, q\right) = H\left(\frac{p}{\alpha(p,q)}, q\right).$$

Then, by the uniqueness of α, we have

$$\frac{\lambda}{\alpha(\lambda p, q)} = \frac{1}{\alpha(p,q)} \Rightarrow \alpha(\lambda p, q) = \lambda \alpha(p,q). \tag{2.9}$$

Proposition 2.3. *For every $M > M^*$, the energy hypersurface $H^{-1}(M)$ is C^2 diffeomorphic to $S^{2n-\ell-1} \times \mathbb{R}^\ell$.*

Proof. By (2.8), we have

$$H^{-1}(M) = \alpha^{-1}(1) = \bigcup_{q \in \mathbb{R}^\ell} \{p \in \mathbb{R}^{2n-\ell} \mid \alpha(p,q) = 1\}. \tag{2.10}$$

Define two maps

$$F_1 : H^{-1}(M) \to S^{2n-\ell-1} \times \mathbb{R}^\ell,\ F_1(p,q) := \left(\frac{p}{|p|}, q\right), \tag{2.11}$$

$$F_2 : S^{2n-\ell-1} \times \mathbb{R}^\ell \to H^{-1}(M),\ F_2(p,q) := \left(\frac{p}{\alpha(p,q)}, q\right). \tag{2.12}$$

For any $(p,q) \in H^{-1}(M)$, we have $\alpha(p,q) = 1$ and

$$F_2 \circ F_1(p,q) = F_2\left(\frac{p}{|p|}, q\right) = \left(\frac{\frac{p}{|p|}}{\alpha\left(\frac{p}{|p|}, q\right)}, q\right) = \left(\frac{p}{\alpha(p,q)}, q\right) = (p,q).$$

This proves $F_2 \circ F_1 = id_{H^{-1}(M)}$. Similarly, we have $F_1 \circ F_2 = id_{S^{2n-\ell-1} \times \mathbb{R}^\ell}$. By (i) of Proposition 2.1, we have $H^{-1}(M) \subset \mathbb{R}^{2n}_*$. Since $\alpha \in C^2(\mathbb{R}^{2n}_*, \mathbb{R})$, then F_2 defined by (2.12) is a C^2-diffeomorphism. ■

Example 2.4. Consider the Hamiltonian function $H : \mathbb{R}^4 \to \mathbb{R}$ defined by $H(p_1,p_2,q_1,q_2) := \frac{1}{2}(|p_1|^2 + |p_2|^2) - \cos(q_1) - \cos(q_2)$. For every $M > 2$, the energy hypersurface $H^{-1}(M)$ consists of circles with radius varying periodically from $\sqrt{2(M-2)}$ to $\sqrt{2(M+2)}$.

2.2 Symplectic dilation

In this subsection, we prepare some results for the multiplicity proof in Section 4.

Definition 2.5. **([12])** A ***symplectic dilation*** of a hypersurface S in a symplectic manifold (M, ω) is a vector field $\xi : U \to TM$ defined on an open neighborhood U of S in M which is transverse to S and satisfies

$$L_\xi \omega = \omega. \tag{2.13}$$

A. Weinsten [12] gives the following equivalent characterization of contact type.

Proposition 2.6. *A hypersurface S in (M,ω) is of* ***contact type*** *if and only if there exists a symplectic dilation of S.*

Let $\xi : U\to TM$ be a symplectic dilation of a hypersurface S in (M,ω). For each $x \in U$, there exist $\delta(x) > 0$, a neighborhood $U(x)$ of x in U and a map $\Phi : (-\delta(x), \delta(x)) \times U(x)\to U(x)$ such that

$$\begin{cases} \dfrac{\mathrm{d}}{\mathrm{d}s}\Phi(s,y) = \xi(\Phi(s,y)),\ s \in (-\delta(x), \delta(x)), \\ \Phi(0,y) = y,\ y \in U(x). \end{cases} \tag{2.14}$$

If there exists a constant $\delta := \delta(S) > 0$ such that $\delta(x) \equiv \delta,\ \forall x \in S$, then

$$\Phi : (-\delta, \delta) \times S\to\Phi((-\delta,\delta) \times S) := \mathcal{N}_\delta \tag{2.15}$$

is a diffeomorphism. If ξ is C^N, where $N \in \mathbb{N} \cup \{+\infty\}$, then Φ is C^N.

Definition 2.7. We call Φ a ***symplectic δ-dilation*** of S. The 1-parameter diffeomorphism group

$$\{\phi^s := \Phi(s,\cdot)\}_{|s|<\delta}$$

is also called a symplectic δ-dilation of S.

Let $\{\phi^s\}_{|s|<\delta}$ be a symplectic δ-dilation of a hypersurface S in (M,ω). A subsequent property of (2.13) is

$$(\phi^s)^*\omega = e^s\omega,\ \forall s \in (-\delta, \delta), \tag{2.16}$$

which is crucial in the proof of (ii) of Proposition 2.8 below. We refer to page 122 of [8] for its proof. The following map

$$\phi^s : \mathcal{N}_{\frac{1}{3}\delta}\to\phi^s(\mathcal{N}_{\frac{1}{3}\delta}),\ \forall s \in (-\frac{2}{3}\delta, \frac{2}{3}\delta), \tag{2.17}$$

is a diffeomorphism. By (2.15) and Definition 2.7, we have

$$\mathcal{N}_\delta = \bigcup_{|s|<\delta} S_s,\ \text{where } S_s := \phi^s(S). \tag{2.18}$$

Choose $\hat{H} \in C^2(M,\mathbb{R})$ such that

$$\hat{H}(x) := f(s),\ \forall s \in (-\frac{2}{3}\delta, \frac{2}{3}\delta),\ x \in S_s, \tag{2.19}$$

where $f:\mathbb{R}\to\mathbb{R}$ is a C^2 auxiliary function and satisfies

$$\begin{cases} f(s)\equiv \max_{s\in\mathbb{R}} f, & s\geqslant \frac{1}{3}\delta, \\ f'(s)>0, & |s|<\frac{1}{3}\delta. \end{cases} \tag{2.20}$$

See Fig 2. By (2.15), the smoothness of $\hat{H}$ follows from that of f.

Proposition 2.8. *Let S be a contact hypersurface in a symplectic manifold (M,ω), $\xi : U\to TM$ be a symplectic dilation of S which generates $\{\phi^s\}_{|s|<\delta}$, $\hat{H}$ be defined by (2.19) and $X_{\hat{H}}$ be the Hamiltonian vector field of $\hat{H}$. The following holds*

(i) For each $s\in(-\frac{1}{3}\delta,\frac{1}{3}\delta)$ and $x\in S_s$, we have

$$\hat{H}'(x)\cdot\xi(x)=f'(s)>0 \tag{2.21}$$

and S_s is a regular energy hypersurface of $\hat{H}$.

(ii) For $s_1,s_2\in(-\frac{1}{3}\delta,\frac{1}{3}\delta)$, if $x_1(t)$ lies on D_{s_1} and solves $\dot{x}_1(t)=X_{\hat{H}}(x_1(t))$, then

$$x_2(t):=\phi^{s_2-s_1}(x_1(t))$$

lies on S_{s_2} and solves

$$\dot{x}_2(t)=e^{s_2-s_1}\frac{f'(s_1)}{f'(s_2)}X_{\hat{H}}(x_2(t)). \tag{2.22}$$

(iii) If F is another function possessing S as a regular energy surface, then

$$X_F(x)=(F'(x)\cdot\xi(x))X_{\hat{H}}(x),\ \forall x\in S. \tag{2.23}$$

Proof. For (i), the conclusion follows from differentiating (2.19) with respect to s.

For (ii), since $\{\phi^s\}$ is generated by ξ, $\hat{H}$ and $\hat{H}\circ\phi^{s_1-s_2}$ are constant on S_{s_2} and regular near S_{s_2}, by (2.21), we have

$$(\hat{H}\circ\phi^{s_1-s_2})'(x)=\frac{f'(s_1)}{f'(s_2)}\hat{H}'(x),\ \forall s_1,s_2\in(-\frac{1}{3}\delta,\frac{1}{3}\delta),\ x\in S_{s_2}. \tag{2.24}$$

Note that

$$d\hat{H}(x_1(t))=d(\hat{H}\circ\phi^{s_1-s_2})(x_2(t))d(\phi^{s_2-s_1})(x_1(t)). \tag{2.25}$$

Then, for any vector field ζ on $\mathcal{N}_{\frac{1}{3}\delta}$, we have

$$\begin{aligned} &\omega\left(\frac{\mathrm{d}}{\mathrm{dt}}x_2(t), d(\phi^{s_2-s_1})(x_1(t))\zeta(x_1(t))\right)\\ =&\omega\left(d(\phi^{s_2-s_1})(x_1(t))\frac{\mathrm{d}}{\mathrm{dt}}x_1(t), d(\phi^{s_2-s_1})(x_1(t))\zeta(x_1(t))\right)\\ =&e^{s_2-s_1}\omega\left(\frac{\mathrm{d}}{\mathrm{dt}}x_1(t), \zeta(x_1(t))\right) = -e^{s_2-s_1}d\hat{H}(x_1(t))\zeta(x_1(t))\\ =&-e^{s_2-s_1}d(\hat{H}\circ\phi^{s_1-s_2})(x_2(t))d(\phi^{s_2-s_1})(x_1(t))\zeta(x_1(t))\\ =&-e^{s_2-s_1}\frac{f'(s_1)}{f'(s_2)}d\hat{H}(x_2(t))d(\phi^{s_2-s_1})(x_1(t))\zeta(x_1(t))\\ =&\omega\left(e^{s_2-s_1}\frac{f'(s_1)}{f'(s_2)}X_{\hat{H}}(x_2(t)), d(\phi^{s_2-s_1})(x_1(t))\zeta(x_1(t))\right). \end{aligned} \tag{2.26}$$

The second equality follows from (2.16). The fourth equality follows from (2.25). The fifth equality follows from (2.24). Since ω is non-degenerate and $d(\phi^{s_2-s_1})(x_1(t)) : T_{x_1(t)}M \to T_{x_2(t)}M$ is isomorphic, by (2.26), we have

$$\frac{\mathrm{d}}{\mathrm{dt}}x_2(t) = e^{s_2-s_1}\frac{f'(s_1)}{f'(s_2)}X_{\hat{H}}(x_2(t)).$$

For (iii), since F and $\hat{H}$ are constant on S and regular near S, then the conclusion follows from (2.21) with $s = 0$. ■

By elementary calculations, we have

Proposition 2.9. *Let $\tilde{M}$ be a covering space of M with covering projection $\pi : \tilde{M} \to M$. The following holds.*

(i) If ω is a symplectic form on M, then $\tilde{\omega}$ defined by

$$\tilde{\omega} := \pi^*\omega \tag{2.27}$$

is a symplectic form on $\tilde{M}$.

(ii) If $\xi : U \subset M \to TM$ is a symplectic dilation of a hypersurface S in (M, ω), then $\tilde{\xi}$ defined by

$$\tilde{\xi}(\tilde{x}) := d\pi(\tilde{x})^{-1}\xi(\pi(\tilde{x})),\ \forall \tilde{x} \in \tilde{U} := \pi^{-1}(U), \tag{2.28}$$

is a symplectic dilation of $\tilde{S} := \pi^{-1}(S)$.

(iii) If $\{\phi^s\}_{|s|<\delta}$ is a symplectic δ-dilation of a hypersurface S in (M,ω), there exists a symplectic δ-dilation $\{\tilde{\phi}^s\}_{|s|<\delta}$ of $\tilde{S}$ such that

$$\pi\circ\tilde{\phi}^s(\tilde{x})=\phi^s\circ\pi(\tilde{x}),\ \forall s\in(-\delta,\delta),\ \tilde{x}\in\tilde{S}. \tag{2.29}$$

2.3 Hamiltonian function $\hat{H}$ for the multiplicity proof

In this subsection, we list some properties of $\bar{H}$ given by P. Felmer [6] for the fixed energy problem (1.4). For the fixed periodic problem (1.5), we define a new Hamiltonian function $\hat{H}$ similar to (2.19), which possesses $H^{-1}(M)$ as a component of its regular energy hypersurface.

Proposition 2.10. (cf. (1.9)-(1.16) of [6]) *For every $M>M^*$, there exists a function $\bar{H}:\mathbb{R}^{2n}\to\mathbb{R}$ such that*

(i) $\bar{H}$ satisfies (H0), (H1) and

$$(\bar{\mathrm{H}}2)\quad 0<2\bar{H}(x)\leqslant p\cdot\bar{H}_p(x),\ \forall x=(p,q)\in\mathbb{R}^{2n}_*.$$

(ii) There exists constants $a_1,a_2,a_3,a_4>0$ such that

$$a_1|p|^2\leqslant\bar{H}(x)\leqslant a_2|p|^2,\ \forall x=(p,q)\in\mathbb{R}^{2n-\ell}\times\mathbb{R}^{\ell}, \tag{2.30}$$

$$2a_1|p|\leqslant|\bar{H}'(x)|\leqslant a_3(|p|+1),\forall x=(p,q)\in\mathbb{R}^{2n-\ell}\times\mathbb{R}^{\ell}, \tag{2.31}$$

$$|\bar{H}''(x)|\leqslant a_4,\ \forall x=(p,q)\in\mathbb{R}^{2n}_*. \tag{2.32}$$

(iii) $\bar{H}$ is regular near $\bar{H}^{-1}(M)=H^{-1}(M)$ and

$$p\cdot\bar{H}_p(p,q)=2M>0,\ \forall(p,q)\in\bar{H}^{-1}(M). \tag{2.33}$$

Proposition 2.11. *Assume that $H:\mathbb{R}^{2n}\to\mathbb{R}$ satisfies (H0)-(H2) with $\ell=n$. For M^* defined by (1.3) and every $M>M^*$, the energy surface $H^{-1}(M)$ is of contact type in the sense of [12] (see Definition 2.5 and Proposition 2.6).*

Proof. Define a vector field

$$\xi:\mathbb{R}^{2n}_*\to\mathbb{R}^{2n}_*,\ (p,q)\mapsto(p,0),\ \forall(p,q)\in\mathbb{R}^{2n}_*. \tag{2.34}$$

By (i) of Proposition 2.10, we have

$$\bar{H}'(e^s p,q)\cdot\xi(e^s p,q)=\bar{H}_p(e^s p,q)\cdot e^s p\geqslant 2\bar{H}(e^s p,q)>0,\ \forall s\in\mathbb{R},\ (p,q)\in\mathbb{R}^{2n}_*.$$

This proves that ξ is a symplectic dilation of $\bar{H}^{-1}(M)=H^{-1}(M)$. ■

Remark 2.12. The vector field ξ defined by (2.34) generates the flow

$$\Phi:\mathbb{R}\times\mathbb{R}^{2n}_*\to\mathbb{R}^{2n}_*,\ (s,p,q)\mapsto(e^s p,q),\ \forall s\in\mathbb{R},\ (p,q)\in\mathbb{R}^{2n}_*,\tag{2.35}$$

which induces diffeomorphisms

$$\Phi:\mathbb{R}\times\bar{H}^{-1}(M)\to\mathbb{R}^{2n}_*\tag{2.36}$$

and

$$\phi^s:\mathbb{R}^{2n}_*\to\mathbb{R}^{2n}_*,\ (p,q)\mapsto(e^s p,q).\tag{2.37}$$

For H satisfying (H0)-(H2), assume that $H^{-1}(M)$ is contact. By Proposition 2.1, the hypersurface $H^{-1}(M)/\mathbb{Z}^\ell\subset\mathbb{R}^{2n-\ell}\times\mathbb{R}^\ell/\mathbb{Z}^\ell$ is compact. By Propositions 2.9-2.11, there exist a constant $\delta>0$ sufficiently small, two symplectic δ-dilations $\{\phi^s\}$ and $\{\hat{\phi}^s\}$ of $H^{-1}(M)$ and $H^{-1}(M)/\mathbb{Z}^\ell$ respectively such that (2.29) holds. Set

$$\mathcal{N}_\delta:=\bigcup_{|s|<\delta}D_s,\ D_s:=\phi^s(D_0),\ D_0:=H^{-1}(M).\tag{2.38}$$

Proposition 2.13. *For every $s\in(-\delta,\delta)$, we have*

$$D_s=D_s+\{0\}\times\mathbb{Z}^\ell\tag{2.39}$$

and

$$\phi^{-s}(x+(0,m))=\phi^{-s}(x)+(0,m),\ x\in D_s,\ m\in\mathbb{Z}^\ell.\tag{2.40}$$

Proof. For any $s\in(-\delta,\delta),\ y\in D_0,\ m\in\mathbb{Z}^\ell$, by (2.3) and (2.29), we have $y+(0,m)\in D_0$ and

$$\pi\circ\phi^s(y+(0,m))=\hat{\phi}^s\circ\pi(y+(0,m))=\hat{\phi}^s\circ\pi(y)=\pi\circ\phi^s(y).$$

Since $\phi^s(y+(0,m))-\phi^s(y)$ is continuous in s, there exists $m'\in\mathbb{Z}^\ell$ independent on s such that

$$(0,m')=\phi^s(y+(0,m))-\phi^s(y)=\phi^0(y+(0,m))-\phi^0(y)=y+(0,m)-y=(0,m).$$

Then $m' = m$ and

$$\phi^s(y + (0, m)) = \phi^s(y) + (0, m),\ \forall s \in (-\delta, \delta),\ y \in D_0,\ m \in \mathbb{Z}^\ell.$$

This proves (2.39) and (2.40). ■

Define auxiliary functions $f, g : \mathbb{R} \to \mathbb{R}$ by

$$f(s) := \begin{cases} 0 & , \quad s \leqslant -\frac{1}{3}\delta, \\ \dfrac{1}{(\frac{1}{3}\delta)^2}(s + \dfrac{1}{3}\delta)^3 + \dfrac{-\frac{1}{2}}{(\frac{1}{3}\delta)^3}(s + \dfrac{1}{3}\delta)^4 & , \quad -\frac{1}{3}\delta \leqslant s \leqslant 0, \\ \dfrac{1}{3}\delta + \dfrac{1}{(\frac{1}{3}\delta)^2}(s - \dfrac{1}{3}\delta)^3 + \dfrac{\frac{1}{2}}{(\frac{1}{3}\delta)^3}(s - \dfrac{1}{3}\delta)^4 & , \quad 0 \leqslant s \leqslant \frac{1}{3}\delta, \\ \dfrac{1}{3}\delta & , \quad s \geqslant \frac{1}{3}\delta, \end{cases} \tag{2.41}$$

$$g(s) := \frac{f'(s)}{e^s}. \tag{2.42}$$

By direct calculation, we have

Proposition 2.14.

(i) f is C^2 and satisfies (2.20).

(ii) g is C^1 and satisfies

$$g(s) \begin{cases} = 0, & |s| \geqslant \dfrac{1}{3}\delta, \\ > 0, & |s| < \dfrac{1}{3}\delta, \end{cases} \text{ and } \begin{cases} \text{increases strictly in } s \in [-\dfrac{1}{3}\delta, -\dfrac{1}{3}\delta + \delta^+], \\ \text{decreases strictly in } s \in [-\dfrac{1}{3}\delta + \delta^+, \dfrac{1}{3}\delta], \end{cases} \tag{2.43}$$

where

$$\delta^+ := \frac{3 + \frac{1}{2}\delta - \sqrt{(3 + \frac{1}{2}\delta)^2 - 4\delta}}{2}. \tag{2.44}$$

See Fig 3.

Proposition 2.15. (Periodic extension) *For the vector $k \in \mathbb{Z}^\ell \setminus \{0\}$ in Theorem 1.1, there exist an $N_0 \in \mathbb{N}$ sufficiently large and a function $\hat{H} \in C^2(\mathbb{R}^{2n}, \mathbb{R})$ such that*

(i) $\hat{H}$ is periodic in all components in the sense of

$$\hat{H}(x + (m', m)) = \hat{H}(x),\ \forall x \in \mathbb{R}^{2n}, (m', m) \in N_0\mathbb{Z}^{2n-\ell} \times \mathbb{Z}^\ell, \tag{2.45}$$

and possesses bounded gradient and Hessian, i.e.,

$$0 < M_1 := \max_{x\in\mathbb{R}^{2n}} |\hat{H}'(x)|,\ M_2 := \max_{x\in\mathbb{R}^{2n}} |\hat{H}''(x)| < +\infty. \tag{2.46}$$

(ii) For each $b \in (0, \frac{1}{3}\delta)$, there exists some $\delta_b \in (-\frac{1}{3}\delta, \frac{1}{3}\delta)$ such that

$$\hat{H}^{-1}(b) = D_{\delta_b} + N_0\mathbb{Z}^{2n-\ell} \times \{0\}, \tag{2.47}$$

where D_s is defined by (2.38). Moreover,

$$\hat{H}'(x) \neq 0 \quad \textit{iff} \quad 0 < \hat{H}(x) < \frac{1}{3}\delta. \tag{2.48}$$

(iii) If $(\bar{x}, \bar{T})$ solves (1.4) with $\bar{T} \neq 0$, then

$$\hat{x}(s) := \bar{x}(t(s)) + t(s)(0,k) - s(0,k) \tag{2.49}$$

solves (1.5) with $\hat{T}$ defined by

$$\hat{T} := \bar{T}\int_0^1 \bar{H}'(\bar{x}(t) + t(0,k)) \cdot \xi(\bar{x}(t) + t(0,k))\mathrm{d}t \neq 0, \tag{2.50}$$

where ξ denotes the symplectic dilation of D_0 which generates ϕ^s in (2.38) and $t(s)$ is defined by

$$\begin{cases} \dfrac{\mathrm{d}t}{\mathrm{d}s} = \hat{T}(\bar{T}\bar{H}'(\bar{x}(t) + t(0,k)) \cdot \xi(\bar{x}(t) + t(0,k)))^{-1} \neq 0, \\ t(0) = 0, \end{cases} \tag{2.51}$$

(iv) If $\hat{x}$ solves (1.5), then $\hat{H}(\hat{x}(t) + t(0,k)) \equiv b \in (0, \frac{1}{3}\delta)$. Moreover,

(iv.1) Each element of

$$[\hat{x}] := \hat{x} + N_0\mathbb{Z}^{2n-\ell} \times \mathbb{Z}^\ell \tag{2.52}$$

solves (1.5).

(iv.2) There exist $\delta_b \in (-\frac{1}{3}\delta, \frac{1}{3}\delta)$ and $x \in [\hat{x}]$ solving (1.5) such that $x(t) + t(0,k)$ lies on D_{δ_b}. Then

$$w(t) := \phi^{-\delta_b}(x(t) + t(0,k)) \in D_0,\ \forall t \in \mathbb{R}. \tag{2.53}$$

(iv.3) $\pi \circ w$ is a non-constant 1-periodic solution of

$$\begin{cases} \dfrac{\mathrm{d}}{\mathrm{d}t}\pi \circ w(t) = g(\delta_b)\hat{T}X_{\hat{H}_\pi}(\pi \circ w(t)), \\ \pi \circ w(1) = \pi \circ w(0), \end{cases} \tag{2.54}$$

where π *denotes the covering projection* $\mathbb{R}^{2n} \to \mathbb{R}^{2n-\ell} \times \mathbb{R}^\ell/\mathbb{Z}^\ell$*,* $\hat{H}_\pi : \mathbb{R}^{2n-\ell} \times \mathbb{R}^\ell/\mathbb{Z}^\ell \to \mathbb{R}$ *is a Hamiltonian function defined by*

$$\hat{H}_\pi \circ \pi := \hat{H}, \tag{2.55}$$

$X_{\hat{H}_\pi}$ *denotes the Hamiltonian vector field of* $\hat{H}_\pi$ *on* $(\mathbb{R}^{2n-\ell} \times \mathbb{R}^\ell/\mathbb{Z}^\ell, \omega_\pi)$ *with* ω_π *determined by*

$$\omega = \pi^* \omega_\pi, \tag{2.56}$$

and ω *denotes the standard symplectic form on* $\mathbb{R}^{2n}$*.*

(iv.4) Denote by $\tilde{\mathscr{P}}$ *the set of periods of* $\pi \circ w$*. There exists* $n(x) \in \mathbb{N}$ *such that*

$$\tilde{\mathscr{P}} = \frac{1}{n(x)}\mathbb{Z}. \tag{2.57}$$

(iv.5)

$$z(s) := w\left(t\left(\frac{s}{T}\right)\right) \tag{2.58}$$

solves (1.1) with T *defined by*

$$T := g(\delta_b)\hat{T}\int_0^1 \frac{\mathrm{d}t}{H'(w(t)) \cdot \xi(w(t))} \neq 0, \tag{2.59}$$

where $\hat{s} := \frac{s}{T}$ *and* $t(\hat{s})$ *is defined by*

$$\begin{cases} \dfrac{\mathrm{d}t}{\mathrm{d}\hat{s}} = \dfrac{TH'(w(t)) \cdot \xi(w(t))}{g(\delta_b)\hat{T}} \neq 0, \\ t(0) = 0. \end{cases} \tag{2.60}$$

Proof. We carry out the proof in three steps.

Step 1. Definition of $\hat{H}$.

Define a map Φ_1 such that the following diagram commutes, i.e.,

$$\Phi_1 := \Phi \circ (id \times F_2) \circ (F_3 \times id), \tag{2.61}$$

$$\begin{tikzcd}
(\mathrm{int} B^{2n-\ell}_{e^\delta}(0) \setminus B^{2n-\ell}_{e^{-\delta}}(0)) \times \mathbb{R}^\ell \arrow[r, "\Phi_1"] \arrow[d, "F_3 \times id"] & \mathcal{N}_\delta \arrow[r, "\breve{H}"] & \mathbb{R} \\
(-\delta, \delta) \times S^{2n-\ell-1} \times \mathbb{R}^\ell \arrow[r, "id \times F_2"] & (-\delta, \delta) \times H^{-1}(M) \arrow[u, "\Phi"'] &
\end{tikzcd} \tag{2.62}$$

$$\begin{tikzcd}
(p,q) \arrow[r, mapsto] \arrow[d, mapsto] & \Phi(\ln|p|, F_2(\frac{p}{|p|}, q)) \arrow[r, mapsto] & f(\ln|p|) \\
(\ln|p|, \frac{p}{|p|}, q) \arrow[r, mapsto] & (\ln|p|, \frac{p}{\alpha(p,q)}, q) \arrow[u, mapsto] &
\end{tikzcd}$$

where Φ is the diffeomorphism (2.15) with $S = H^{-1}(M)$, $F_2 : S^{2n-\ell-1} \times \mathbb{R}^{\ell} \to H^{-1}(M)$ is the C^2-diffeomorphism defined by (2.12), $F_3 : \mathrm{int}B_{e^{\delta}}^{2n-\ell}(0) \setminus B_{e^{-\delta}}^{2n-\ell}(0) \to (-\delta, \delta) \times S^{2n-\ell-1}$ is defined by

$$F_3(p) := (\ln|p|, \frac{p}{|p|}),\ \forall p \in B_{e^{\delta}}^{2n-\ell}(0) \setminus \bar{B}_{e^{-\delta}}^{2n-\ell}(0), \tag{2.63}$$

and

$$B_{\rho}^{2n-\ell}(0) := \{p \in \mathbb{R}^{2n-\ell} \mid |p| \leqslant \rho\},\ \mathrm{int}B_{\rho}^{2n-\ell}(0) := \{p \in \mathbb{R}^{2n-\ell} \mid |p| < \rho\},\ \rho > 0. \tag{2.64}$$

Then Φ_1 is a diffeomorphism. The C^2-smoothness of Φ_1 follows from that of $F_3,\ F_2$ and Φ. To continue our study, we define an auxiliary function $\breve{H} : \mathbb{R}^{2n} \to \mathbb{R}$ in two cases according to the value of ℓ in (H1).

Case 1. $1 \leqslant \ell \leqslant 2n-2$. By (2.61) and (2.62) with δ replaced by any $\delta' \in (0, \delta)$, the set $\mathbb{R}^{2n} \setminus \mathcal{N}_{\delta'}$ has two components $V_{\delta'}$ and $W_{\delta'}$ such that $V_{\delta'}/\mathbb{Z}^{\ell}$ is compact and $\{0\} \times \mathbb{R}^{\ell}$ is contained in the interior of $V_{\delta'}$. Choose f defined by (2.41) and define a new Hamiltonian function $\breve{H}$ by

$$\breve{H}(x) := \begin{cases} 0, & x \in V_{\frac{1}{3}\delta}, \\ f(s), & x \in D_s,\ |s| < \frac{2}{3}\delta, \\ \frac{1}{3}\delta, & x \in W_{\frac{1}{3}\delta}, \end{cases} \tag{2.65}$$

Case 2. $\ell = 2n-1$. By (2.65) and (2.66) with δ replaced by any $\delta' \in (0, \delta)$, the set $\mathbb{R}^{2n} \setminus \mathcal{N}_{\delta'}$ has three components $V_{\delta'}$, $W_{\delta'}^{\pm}$ such that $V_{\delta'}/\mathbb{Z}^{\ell}$ is compact and $\{0\} \times \mathbb{R}^{\ell}$ is contained in the interior of $V_{\delta'}$. We can also define $\breve{H}$ similar to (2.65) by

$$\breve{H}(x) := \begin{cases} 0, & x \in V_{\frac{1}{3}\delta}, \\ f(s), & x \in D_s,\ |s| < \frac{2}{3}\delta, \\ \frac{1}{3}\delta, & x \in W_{\frac{1}{3}\delta}^{-} \cup W_{\frac{1}{3}\delta}^{+}. \end{cases} \tag{2.66}$$

It follows from (2.1) that

$$D_0 = H^{-1}(M) \subset B^{2n-\ell}_{r''}(0) \times \mathbb{R}^\ell, \tag{2.67}$$

Since $\delta > 0$ is sufficiently small, by (2.1), (2.38) and (2.67), there exists

$$r_\delta \in \mathbb{N} \cap (r'', \infty) \tag{2.68}$$

such that

$$\mathcal{N}_\delta \subset B^{2n-\ell}_{r_\delta}(0) \times \mathbb{R}^\ell \subset \mathtt{int}\mathcal{O}_\delta \times \mathbb{R}^\ell, \tag{2.69}$$

where the closed set $\mathcal{O}_\delta$ is defined by

$$\mathcal{O}_\delta := \{p = (p_1, p_2, \cdots, p_{2n-\ell}) \in \mathbb{R}^{2n-\ell} \mid |p_i| \leqslant r_\delta + 1,\ 1 \leqslant i \leqslant 2n-\ell\} \tag{2.70}$$

and int$\mathcal{O}_\delta$ denotes the interior of $\mathcal{O}_\delta$. Then $\breve{H}$ defined by (2.65) or (2.66) equals to its maximum $\frac{1}{3}\delta$ on $\mathbb{R}^{2n} \setminus (\mathcal{O}_\delta \times \mathbb{R}^\ell)$. Choose

$$N_0 := 2(r_\delta + 1). \tag{2.71}$$

Then we have

$$\mathbb{R}^{2n-\ell} = \bigcup_{m' \in \mathbb{Z}^{2n-\ell}} (\mathcal{O}_\delta + N_0 m') := \mathcal{O}_\delta + N_0 \mathbb{Z}^{2n-\ell} \tag{2.72}$$

and

$$\begin{aligned} (\mathrm{int}\mathcal{O}_{\delta_b} \times \mathbb{R}^\ell + (m_1', 0)) \cap (\mathrm{int}\mathcal{O}_{\delta_b} \times \mathbb{R}^\ell + (m_2', 0)) = \emptyset, \\ \forall m_1', m_2' \in N_0\mathbb{Z}^{2n-\ell}, m_1' \neq m_2'. \end{aligned} \tag{2.73}$$

Using $\breve{H}$, we can define a new Hamiltonian function $\hat{H} : \mathbb{R}^{2n} \to \mathbb{R}$ by

$$\hat{H}(p + N_0 m', q) = \breve{H}(p, q),\ \forall p \in \mathcal{O}_\delta,\ q \in \mathbb{R}^\ell,\ m' \in \mathbb{Z}^{2n-\ell}. \tag{2.74}$$

It follows from (2.72) and (2.73) that $\hat{H}$ is well-defined. Then $\hat{H}$ satisfies (i)-(ii). Here, we define

$$\breve{F} : (\mathrm{int}B^{2n-\ell}_{e^\delta}(0) \setminus B^{2n-\ell}_{e^{-\delta}}(0)) \times \mathbb{R}^\ell \to \mathbb{R},\ (p, q) \mapsto f(\ln|p|).$$

Then in the diagram (2.62) we have $\breve{F} = \breve{H} \circ \Phi_1$. Since f and Φ_1 are C^2, then $\breve{F}$ and $\breve{H}$ are C^2. Hence $\hat{H}$ is C^2. The periodicity of $\hat{H}$ in q follows from (2.39). The most new important property of $\hat{H}$ is its periodicity in p.

Step 2. Proof of (iii).

Since ξ is a symplectic dilation of D_0, $\bar{H}$ is regular near $\bar{H}^{-1}(M) = H^{-1}(M) = D_0$ and $\bar{H}(\bar{x}(t) + t(0,k)) \equiv M$, then ξ is transverse to D_0 and

$$\bar{H}'(\bar{x}(t) + t(0,k)) \cdot \xi(\bar{x}(t) + t(0,k)) \neq 0. \tag{2.75}$$

It follows from $\bar{T} \neq 0$ and (2.75) that $\hat{T} \neq 0$ and (2.51) is well-defined. By (2.23) with F replaced by $\bar{H}$, we have

$$X_{\bar{H}}(x) = (\bar{H}'(x) \cdot \xi(x)) X_{\hat{H}}(x), \ \forall x \in D_0. \tag{2.76}$$

Since $\bar{H}(\bar{x}(t) + t(0,k)) \equiv M$, we have

$$\hat{x}(s) + s(0,k) = \bar{x}(t(s)) + t(s)(0,k) \in D_0, \ \forall s \in \mathbb{R}. \tag{2.77}$$

Then

$$\begin{aligned}
\frac{\mathrm{d}}{\mathrm{d}s}(\hat{x}(s) + s(0,k)) &= \frac{\mathrm{d}t}{\mathrm{d}s}\frac{\mathrm{d}}{\mathrm{d}t}(\bar{x}(t) + t(0,k)) = \frac{\mathrm{d}t}{\mathrm{d}s}\bar{T} X_{\bar{H}}(\bar{x}(t) + t(0,k)) \\
&= \frac{\mathrm{d}t}{\mathrm{d}s}\bar{T}\big(\bar{H}'(\bar{x}(t) + t(0,k)) \cdot \xi(\bar{x}(t) + t(0,k))\big) X_{\hat{H}}(\bar{x}(t) + t(0,k)) \\
&= \hat{T} X_{\hat{H}}(\hat{x}(s) + s(0,k)).
\end{aligned}$$

The third equality follows from (2.76) and (2.77). The last equality follows from (2.51). By (2.51), the inverse function $s(t)$ of $t(s)$ is given by

$$\begin{cases} \dfrac{\mathrm{d}s}{\mathrm{d}t} = \dfrac{\bar{T}}{\hat{T}}\bar{H}'(\bar{x}(t) + t(0,k)) \cdot \xi(\bar{x}(t) + t(0,k)), \\ s(0) = 0. \end{cases}$$

We obtain $t(1) = 1$ from

$$s(1) = \frac{\bar{T}}{\hat{T}} \int_0^1 \bar{H}'(\bar{x}(t) + t(0,k)) \cdot \xi(\bar{x}(t) + t(0,k)) \mathrm{d}t = 1.$$

Then $\hat{x}(1) = \bar{x}(1) = \bar{x}(0) = \hat{x}(0)$ and $\hat{x}$ solves (1.5).

Step 3. Proof of (iv).

Set

$$\hat{\gamma}(t) := \hat{x}(t) + t(0,k).$$

Since $\hat{x}$ solves (1.5) and $k \neq 0$, there exists a constant b such that $\hat{H}(\hat{\gamma}(t)) \equiv b$ and $\hat{H}' \neq 0$ on $\hat{\gamma}(\mathbb{R})$. By (2.48), we have $b \in (0, \frac{1}{3}\delta)$.

For (iv.1), each element $\tilde{x} \in [\hat{x}]$ can be written as $\tilde{x} = \hat{x} + (m', m)$, where $m' \in N_0\mathbb{Z}^{2n-\ell}$, $m \in \mathbb{Z}^{\ell}$. Since $\hat{x}$ solves (1.5), it follows from (2.45) that $\tilde{x} = \hat{x} + (m', m)$ solves (1.5).

For (iv.2), since $\hat{H}(\hat{\gamma}(t)) \equiv b \in (0, \frac{1}{3}\delta)$, by (iii), there exists $\delta_b \in (-\frac{1}{3}\delta, \frac{1}{3}\delta)$ such that (2.47) holds. Then

$$\hat{\gamma}(\mathbb{R}) \subset D_{\delta_b} + N_0\mathbb{Z}^{2n-\ell} \times \{0\}. \tag{2.78}$$

By (2.39) and (2.69), we have

$$D_{\delta_b} \subset \mathrm{int}\mathcal{O}_\delta \times \mathbb{R}^\ell.$$

By (2.73), we have

$$\begin{aligned}
&(D_{\delta_b} + (m_1', 0)) \cap (D_{\delta_b} + (m_2', 0)) \\
\subset &(\mathrm{int}\mathcal{O}_\delta \times \mathbb{R}^\ell + (m_1', 0)) \cap (\mathrm{int}\mathcal{O}_\delta \times \mathbb{R}^\ell + (m_2', 0)) \\
= &\emptyset, \ \forall m_1', m_2' \in N_0\mathbb{Z}^{2n-\ell}, m_1' \neq m_2'.
\end{aligned} \tag{2.79}$$

Since $\hat{\gamma}(\mathbb{R})$ is connected, by (2.78) and (2.79), there exists a unique $m_b' \in N_0\mathbb{Z}^{2n-\ell}$ such that the orbit $\hat{\gamma}$ lies on $D_{\delta_b} + (m_b', 0)$. Then $x(t) + t(0, k) = \hat{\gamma}(t) - (m_b', 0)$ lies on D_{δ_b}.

For (iv.3), the conclusion follows from (2.22), (2.42) and elementary calculations.

For (iv.4), since $\pi \circ w$ is a non-constant 1-periodic orbit, then $\tilde{\mathscr{P}}$ is well-defined and there exists $n(x) \in \mathbb{N}$ such that (2.57) holds.

For (iv.5), since $x(t) + t(0, k)$ lies on D_{δ_b}, then $w(t)$ lies on D_0. Similarly to (2.75), we have

$$H'(w(t)) \cdot \xi(w(t)) \neq 0. \tag{2.80}$$

Then T determined by (2.59) is well-defined. By $|\delta_b| < \frac{1}{3}\delta$ and Proposition 2.14, we have $g(\delta_b) > 0$. Since $\hat{T} \neq 0$, by (2.21), we have $T \neq 0$. Since x solves (1.5) and $x(t) + t(0, k)$ lies on D_{δ_b}, by (2.22) with $s_1 = \delta_b, s_2 = 0, x_1(t) = x(t), x_2(t) + t(0, k) = w(t)$, we have

$$\frac{\mathrm{d}}{\mathrm{d}t} w(t) = e^{-\delta_b} f'(\delta_b)\hat{T} X_{\hat{H}}(w(t)) = g(\delta_b)\hat{T} X_{\hat{H}}(w(t)). \tag{2.81}$$

The second equality follows from (2.42). Similarly to (2.76), we have

$$X_H(x) = (H'(x) \cdot \xi(x)) X_{\hat{H}}(x), \ \forall x \in D_0. \tag{2.82}$$

It follows from $g(\delta_b), \hat{T}, T \neq 0$ and (2.80) that (2.60) is well-defined. Since $z(s) = w(t)$ lies one D_0, we have

$$\begin{aligned}
\frac{\mathrm{d}}{\mathrm{d}s} z(s) &= \frac{\mathrm{d}\hat{s}}{\mathrm{d}s} \frac{\mathrm{d}t}{\mathrm{d}\hat{s}} \frac{\mathrm{d}}{\mathrm{d}t} w(t) \\
&= \frac{1}{T} \frac{\mathrm{d}t}{\mathrm{d}\hat{s}} g(\delta_b) \hat{T} X_{\hat{H}}(w(t)) \\
&= \frac{1}{T} \frac{\mathrm{d}t}{\mathrm{d}\hat{s}} g(\delta_b) \hat{T} \frac{X_H(z(s))}{H'(w(t)) \cdot \xi(w(t))} \\
&= X_H(z(s)).
\end{aligned}$$

The fourth equality follows from (2.81). The fifth equality follows from (2.82). The last equality follows from (2.60). By (2.60), the inverse function $\hat{s}(t)$ of $t(\hat{s})$ is given by

$$\begin{cases}
\dfrac{\mathrm{d}\hat{s}}{\mathrm{d}t} = \dfrac{g(\delta_b)\hat{T}}{T H'(w(t)) \cdot \xi(w(t))}, \\
\hat{s}(0) = 0.
\end{cases}$$

We obtain $t(1) = 1$ from

$$\hat{s}(1) = \frac{g(\delta_b)\hat{T}}{T} \int_0^1 \frac{\mathrm{d}\tau}{H'(w(\tau)) \cdot \xi(w(\tau))} = 1.$$

Then

$$\begin{aligned}
z(T) - z(0) &= w(1) - w(0) \\
&= \phi^{-\delta_b}(x(1) + (0,k)) - \phi^{-\delta_b}(x(0)) \\
&= \phi^{-\delta_b}(x(0) + (0,k)) - \phi^{-\delta_b}(x(0)) \\
&= (0,k).
\end{aligned}$$

The third equality follows from $x(1) = x(0)$. The last equality follows from $x(0) \in D_{\delta_b}$ and (2.40) with $s = \delta_b$. ■

Remark 2.16. Note that the Hamiltonian function $\breve{H}$ defined by (2.65) or (2.66) also satisfies Proposition 2.15 except that it is not periodic in p. Similar results of [4] listed in Section 3 are valid for $\breve{H}$. However, we can only obtain $n+1$ solutions of (1.5) with $\hat{H}$ replaced by $\breve{H}$. Then, in section 4.1, we can only obtain ${}^{\#}\tilde{\mathcal{P}} \geqslant \frac{n+1}{2}$. Therefore, periodic extension of $\breve{H}$ in p enable us to obtain more solutions of (1.5) and (1.1).

Proposition 2.17. *Assume that H satisfies (H0)-(H2) with $\ell = n$. There exists a solution of (1.5).*

Proof. By the proof of Theorem 0.1 of [6], there exists a solution $(\bar{x}, \bar{T})$ of (1.4). Then by (iv) of Proposition 2.15 there exists a solution of (1.5). ■

3 Saddle point reduction

For the Hamiltonian function $\hat{H} : \mathbb{R}^{2n} \to \mathbb{R}$ and the number $\hat{T}$ given by Proposition 2.15, define two functionals $\mathcal{A}, \mathcal{B} : E = H^1(S^1, \mathbb{R}^{2n}) \to \mathbb{R}$ by

$$\mathcal{A}(x) = \frac{1}{2}\langle Ax, x\rangle_{L^2} - \mathcal{B}(x), \tag{3.1}$$

$$\mathcal{B}(x) = \int_0^1 (\hat{T}\hat{H}(x(t) + t(0,k)) + x \cdot J(0,k))\mathrm{d}t. \tag{3.2}$$

Critical points of $\mathcal{A}$ are exactly solutions to the equation

$$Ax - \mathcal{B}'(x) = 0. \tag{3.3}$$

In this section, we follow the idea of [4] to reduce the functional $\mathcal{A}$ to a functional $G : Z \to \mathbb{R}$ defined on a finite dimensional subspace Z of $L := L^2(\mathbb{S}^1, \mathbb{R}^{2n})$ such that critical points of G are in one-to-one correspondence with that of $\mathcal{A}$ and hence give 1-periodic solutions of (1.5).

Note that the spectrum of the operator $A = -J\frac{\mathrm{d}}{\mathrm{d}t} : E \to L$ is a pure point spectrum and $\sigma(A) = 2\pi\mathbb{Z}$. Let $\{E_\lambda \mid \lambda \in \mathbb{R}\}$ be the spectral resolution of A. In view of (2.46), choose some $C \geqslant 0$ such that $2\pi < 2\hat{T}M_2 + C \notin 2\pi\mathbb{Z} = \sigma(A)$. We can define a orthogonal projection P of L by

$$P := \int_{-2\hat{T}M_2 - C}^{2\hat{T}M_2 + C} \mathrm{d}E_\lambda. \tag{3.4}$$

Set $P^\perp = 1 - P,\ Z = P(L)$ and $Y = P^\perp(L)$. Then $L = Z \oplus Y$ and $\dim Z = 2n + 2d$ for some $d \in \mathbb{N}$, since $2\hat{T}M_2 + C > 2\pi$. Set $A_0 = A|_Y$, $z = Px$ and $y = P^\perp x,\ x \in E$. Equation (3.3) is equivalent to

$$Az - P\mathcal{B}'(z+y) = 0, \tag{3.5}$$

$$Ay - P^{\perp}\mathcal{B}'(z+y) = 0 \Leftrightarrow y = A_0^{-1}P^{\perp}\mathcal{B}'(z+y). \tag{3.6}$$

The right hand side of (3.6) is a contraction operator on L with contraction constant $\frac{1}{2}$. For fixed $z \in Z$, the equation (3.6) has a unique solution $y = y(z) \in E$. Note that x can be uniquely split as $x(z) = z + y(z)$ and equation (3.5) becomes $Az - P\mathcal{B}'(x(z)) = 0$, which is equivalent to $Ax(z) - \mathcal{B}'(x(z)) = 0$. Define

$$G : Z \to \mathbb{R},\, z \mapsto \mathcal{A}(x(z)). \tag{3.7}$$

We have

$$G'(z) = Az - P\mathcal{B}'(x(z)) = Az - \hat{T}P\hat{H}'(x + t(0,k)) - J(0,k). \tag{3.8}$$

In view of (2.45), from the uniqueness of solution of (3.6) follows that

$$y(z + (m', m)) = y(z),\ \forall z \in Z,\ (m', m) \in N_0\mathbb{Z}^{2n-\ell} \times \mathbb{Z}^{\ell}.$$

Hence

$$\begin{aligned} x(z + (m', m)) &= x(z) + (m', m), \quad \text{and} \\ G'(z + (m', m)) &= G'(z),\ \forall z \in Z,\ (m', m) \in N_0\mathbb{Z}^{2n-\ell} \times \mathbb{Z}^{\ell}. \end{aligned} \tag{3.9}$$

Note that if z is a critical point of G, the equivalent class

$$[z] := z + N_0\mathbb{Z}^{2n-\ell} \times \mathbb{Z}^{\ell}$$

is a group of critical points of G with consistent energy. Moreover, the equivalent class

$$[x] := [x(z)] := x(z) + N_0\mathbb{Z}^{2n-\ell} \times \mathbb{Z}^{\ell}$$

is a group of 1-periodic solutions of (1.5) with consistent energy and uniquely determined by the equivalent class $[z]$. Now we write $z \in Z$ as $z = w + \xi$, where $w = \int_0^1 z(t)\mathtt{dt}$. We have $w \in \ker(A)$ and $\xi \in (\ker(A))^{\perp} \cap Z$. Writing $z = (w, \xi)$, we conclude from (3.9) that $G'(z) = G'(w, \xi)$ is a vector field on $T_{\ell}^{2n} \times \mathbb{R}^{2d}$, where T_{ℓ}^{2n} denotes the (non-standard) torus $T_{\ell}^{2n} := \mathbb{R}^{2n}/(N_0\mathbb{Z}^{2n-\ell} \times \mathbb{Z}^{\ell}) \cong \mathbb{R}^{2n}/\mathbb{Z}^{2n} = T^{2n}$. In summary, we have

Proposition 3.1. **(Lemma 1 of [4])** *Rest points of Lipschitz continuous vector field* $G'(z) = G'(w,\xi)$ *on* $T_\ell^{2n} \times \mathbb{R}^{2d}$ *and hence equivalent classes* $[z]$*s, which consist of critical points of the* C^2 *function* $G : Z \to \mathbb{R}$ *defined by (3.7), are in one-to-one correspondence with equivalent classes* $[x]$ *consisting of solutions of (1.5).*

To study rest points of G', we consider the gradient flow

$$\frac{\mathrm{d}}{\mathrm{d}s} z = G'(z) \tag{3.10}$$

on $T_\ell^{2n} \times \mathbb{R}^{2d}$.

Proposition 3.2. **(Corollary of Theorem 5 of [4])** *Denote by* $R(G)$ *the set of rest points of gradient flow (3.10). If* $R(G) \neq \emptyset$*, then* ${}^{\#}R(G) \geqslant 2n+1$.

Remark 3.3. Propositions 3.1-3.2 are basically the same with corresponding results in [4] except that the potential Φ in [4] is periodic in $2n$ directions, while in this section the potential $\mathcal{B}$ defined by (3.2) is only periodic in $2n-1$ directions since $k \neq 0$. However, gradients of Φ and $\mathcal{B}$ are all periodic in $2n$ directions. Hence, the idea of [4] is valid for the multiplicity proof of solutions of (1.5).

Remark 3.4. By Propositions 2.17 and 3.1-3.2, if H satisfies (H0)-(H2) with $\ell = n$, there exist $(x_i, b_i, \delta_{b_i}),\ 1 \leqslant i \leqslant 2n+1$ such that x_i solves (1.5), $b_i \in (0, \frac{1}{3}\delta),\ \delta_{b_i} \in (-\frac{1}{3}\delta, \frac{1}{3}\delta),\ x_i(t) + t(0,k)$ lies on $D_{\delta_{b_i}}$ and x_is are geometrically distinct in the sense of (1.2). In the next section, these $2n+1$ solutions are classified into two categories and give at least $n+1$ geometrically distinct solutions of (1.1).

4 Proof of Theorems

4.1 Proof of Theorem 1.1

Proposition 4.1. *Let* $k_i \in \mathbb{Z}^\ell \setminus \{0\}$ *for* $i \in \{1,2\}$*. Assume that* (x_i, τ_i) *solves (1.5) with* $(k, \hat{T})$ *being replaced by* (k_i, τ_i) *and* $x_i(s) + s(0,k_i)$ *lies on* D_{δ_i} *with* $\delta_i \in (-\frac{1}{3}\delta, \frac{1}{3}\delta)$*. If*

$$w_i(s) := \phi^{-\delta_i}(x_i(s) + s(0,k_i)),\ i = 1,2,$$

are geometrically the same in the sense of

$$\pi \circ w_1(\mathbb{R}) = \pi \circ w_2(\mathbb{R}), \tag{4.1}$$

where π *denotes the covering projection* $\mathbb{R}^{2n} \to \mathbb{R}^{2n-\ell} \times \mathbb{R}^\ell/\mathbb{Z}^\ell$*. Then*

(i) There exists a reparametrization $s : \mathbb{R} \to \mathbb{R},\ t \mapsto s(t)$ *such that*

$$\dot{s}(t) \equiv \frac{g(\delta_1)\tau_1}{g(\delta_2)\tau_2},\ \forall t \in \mathbb{R}, \tag{4.2}$$

$$\pi \circ w_1(t) = \pi \circ w_2(s(t)),\ \forall t \in \mathbb{R}, \tag{4.3}$$

$$k_1 = \dot{s}(0)k_2, \tag{4.4}$$

where $g : \mathbb{R} \to \mathbb{R}$ *is defined by (2.42).*

(ii) There exists $\epsilon \in \{\pm 1\}$ *such that*

$$\frac{g(\delta_1)\tau_1}{g(\delta_2)\tau_2} = \epsilon \frac{|k_1|}{|k_2|}. \tag{4.5}$$

Proof. For (i), without loss of generality, we assume that

$$\pi \circ w_1(0) = \pi \circ w_2(0). \tag{4.6}$$

Since (x_i, τ_i) solves (1.5) with $(k, \hat{T})$ being replaced by (k_i, τ_i) and $x_i(s) + s(0, k_i)$ lies on D_{δ_i} with $\delta_i \in (-\frac{1}{3}\delta, \frac{1}{3}\delta)$, it follows from (v.3) of Proposition 2.15 that $\pi \circ w_i$ solves

$$\begin{cases} \dfrac{\mathrm{d}}{\mathrm{d}t}\pi \circ w_i(t) = g(\delta_i)\tau_i X_{\hat{H}_\pi}(\pi \circ w_i(t)), \\ \pi \circ w_i(1) = \pi \circ w_i(0). \end{cases} \tag{4.7}$$

Define $s : \mathbb{R} \to \mathbb{R}$ by

$$s(t) := \dot{s}(0)t := \frac{g(\delta_1)\tau_1}{g(\delta_2)\tau_2}t,\ \forall t \in \mathbb{R}. \tag{4.8}$$

Set

$$\bar{w}_1(t) := w_2(s(t)). \tag{4.9}$$

Then

$$\begin{aligned}\frac{\mathrm{d}}{\mathrm{d}t}\pi\circ\bar{w}_1(t) &= \frac{\mathrm{d}s}{\mathrm{d}t}\frac{\mathrm{d}}{\mathrm{d}s}\pi\circ w_2(s)\\ &= \dot{s}(t)g(\delta_2)\tau_2 X_{\hat{H}_\pi}(\pi\circ w_2(s))\\ &= g(\delta_1)\tau_1 X_{\hat{H}_\pi}(\pi\circ\bar{w}_1(t)), \qquad (4.10)\\ \pi\circ\bar{w}_1(0) &= \pi\circ w_2(0) = \pi\circ w_1(0). \qquad (4.11)\end{aligned}$$

The second equality follows from (4.7). The third equality follows from (4.8). By (4.7) with $i=1$, (4.10) and (4.11), the basic uniqueness theorem for the initial value problem of ordinary differential equations yields

$$\pi\circ w_1(t) = \pi\circ\bar{w}_1(t),\ \forall t\in\mathbb{R}. \tag{4.12}$$

Then (4.3) follows from (4.9) and (4.12).

By (4.2) and (4.3), we have

$$\begin{aligned}\pi\circ w_2(s+\dot{s}(0)) = \pi\circ w_2(s(t)+\dot{s}(0)) = \pi\circ w_2(\dot{s}(0)(t+1)) = \pi\circ w_2(s(t+1))\\ = \pi\circ w_1(t+1) = \pi\circ w_1(t) = \pi\circ w_2(s(t)) = \pi\circ w_2(s).\end{aligned}$$

This proves that $\dot{s}(0)$ is a period of $\pi\circ w_2$. Since (x_2,τ_2) solves (1.5) with $(k,\hat{T})$ being replaced by (k_2,τ_2) and $x_2(s)+s(0,k_2)$ lies on D_{δ_2} with $\delta_2\in(-\frac{1}{3}\delta,\frac{1}{3}\delta)$, by (v.5) of Proposition 2.15, there exists $m_2\in\mathbb{Z}\setminus\{0\},\ n(x_2)\in\mathbb{N}$ such that

$$\dot{s}(0) = \frac{m_2}{n(x_2)}. \tag{4.13}$$

By (4.2), (4.3) and (4.13), there exists $j\in\mathbb{Z}^\ell$ such that

$$\phi^{-\delta_1}(x_1(t)+t(0,k_1)) = \phi^{-\delta_2}\left(x_2\left(\frac{m_2}{n(x_2)}t\right)+\frac{m_2}{n(x_2)}t(0,k_2)\right)+(0,j). \tag{4.14}$$

Taking the difference of (4.14) with $t = 0$ and $n(x_2)$, we have

$$\begin{aligned}
n(x_2)(0,k_1) &= \phi^{-\delta_1}(x_1(0) + n(x_2)(0,k_1)) - \phi^{-\delta_1}(x_1(0)) \\
&= \phi^{-\delta_1}(x_1(n(x_2)) + n(x_2)(0,k_1)) - \phi^{-\delta_1}(x_1(0)) \\
&= \phi^{-\delta_1}(x_1(t) + t(0,k_1))\Big|_{t=0}^{t=n(x_2)} \\
&= \phi^{-\delta_2}\left(x_2\left(\frac{m_2}{n(x_2)}t\right) + \frac{m_2}{n(x_2)}t(0,k_2)\right)\Big|_{t=0}^{t=n(x_2)} \\
&= \phi^{-\delta_2}(x_2(m_2) + m_2(0,k_2)) - \phi^{-\delta_2}(x_2(0)) \\
&= \phi^{-\delta_2}(x_2(0) + m_2(0,k_2)) - \phi^{-\delta_2}(x_2(0)) \\
&= m_2(0,k_2).
\end{aligned} \tag{4.15}$$

The first and the last equalities follow from $x_i(0) \in D_{\delta_i},\ n(x_2)k_1, m_2k_2 \in \mathbb{Z}^\ell$ and (2.40) with $s_i = \delta_i$ respectively. The second and the fifth equalities follow from $n(x_2) \in \mathbb{N},\ m_2 \in \mathbb{Z}\setminus\{0\}$ and x_i is 1-periodic. Then (4.4) follows from (4.13) and (4.15).

For (ii), the conclusion follows from (4.2) and (4.4). ■

Given $k \in \mathbb{Z}^n \setminus \{0\}$, by Remark 3.4, we obtain at least $2n+1$ solutions of (1.5), denoted by $x_1, x_2, \cdots, x_{2n+1}$, which are distinct in the sense of (1.2). There exist $\delta_i \in (-\frac{1}{3}\delta, \frac{1}{3}\delta), 1 \leqslant i \leqslant 2n+1$ such that

$$w_i(t) := \phi^{-\delta_i}(x_i(t) + t(0,k)) \in D_0,\ \forall t \in \mathbb{R},\ 1 \leqslant i \leqslant 2n+1. \tag{4.16}$$

Set

$$\mathcal{P}_1 := \{w_i \mid \delta_i \in (-\frac{1}{3}\delta, -\frac{1}{3}\delta + \delta^+],\ 1 \leqslant i \leqslant 2n+1\}, \tag{4.17}$$

$$\mathcal{P}_2 := \{w_i \mid \delta_i \in [-\frac{1}{3}\delta + \delta^+, \frac{1}{3}\delta),\ 1 \leqslant i \leqslant 2n+1\}, \tag{4.18}$$

where δ^+ is defined by (2.44). Recall that g defined by (2.42) is positive on $(-\frac{1}{3}\delta, \frac{1}{3}\delta)$, increases strictly on $(-\frac{1}{3}\delta, -\frac{1}{3}\delta + \delta^+]$ and decreases strictly on $[-\frac{1}{3}\delta + \delta^+, \frac{1}{3}\delta)$. We have

Proposition 4.2. *For every $i \in \{1,2\}$, elements of $\mathcal{P}_i$ are pair-wise distinct in the sense of (1.2).*

Proof. We argue by contradiction. Suppose that there exists a pair $w_{i_1},\ w_{i_2}$ belonging to $\mathcal{P}_i$ for some $i \in \{1, 2\}$ such that $\pi \circ w_{i_1}(\mathbb{R}) = \pi \circ w_{i_2}(\mathbb{R})$. By (iii) of Proposition 4.1 with $(k_1, \tau_1) = (k_2, \tau_2) = (k, \hat{T})$, we have $g(\delta_{i_1}) = \epsilon_1 g(\delta_{i_2})$. By Proposition 2.14, we have $g(\delta_{i_1}) = g(\delta_{i_2}) > 0$ and hence $\delta_{i_1} = \delta_{i_2}$. By (4.2), we have $\dot{s}(0) = 1$. By (4.13) and (4.14), we have

$$\begin{aligned} x_{i_1}(t) + t(0, k) &= \phi^{\delta_{i_1}}\left(\phi^{-\delta_{i_2}}(x_{i_2}(t) + t(0, k)) + (0, j)\right) \\ &= \phi^{\delta_{i_1}} \circ \phi^{-\delta_{i_2}}(x_{i_2}(t) + t(0, k)) + (0, j) \\ &= x_{i_2}(t) + t(0, k) + (0, j). \end{aligned} \tag{4.19}$$

The second equality follows from (4.16) and (2.40) with $s = \delta_{i_1}$. The last equality follows from $\delta_1 = \delta_2$. However, the equality (4.19) contradicts the assumption $\pi \circ x_{i_1}(\mathbb{R}) \neq \pi \circ x_{i_2}(\mathbb{R})$. ■

Proof of Theorem 1.1. By Proposition 4.2, we have

$$\begin{aligned} &{}^{\#}\{w_i,\ 1 \leqslant i \leqslant 2n+1 \mid w_i\text{s are geometrically distinct in the sense of (1.2).}\} \\ &\geqslant \max\{{}^{\#}\mathcal{P}_1, {}^{\#}\mathcal{P}_2\} \geqslant \frac{2n+1}{2} > n. \end{aligned}$$

By (iv.5) of Proposition 2.15, we have $\tilde{\mathcal{P}}(M, k) \geqslant n + 1$. ■

4.2 Proof of Theorem 1.2

Since $H^{-1}(M)$ is contact, by (2.65) or (2.66), we obtain a Hamiltonian function $\breve{H} : \mathbb{R}^{2n-\ell} \times (\mathbb{R}^\ell/\mathbb{Z}^\ell) \to \mathbb{R}$ which is bounded in $p \in \mathbb{R}^{2n-\ell}$ and possesses $H^{-1}(M)$ as a regular energy hypersurface. Since $2n - \ell \geqslant 1$, by Theorem 1.2 of [9] and (v.3) of Proposition 2.15, there exists a nonconstant solution (z, T) of (1.1) with some $k_0 \in \mathbb{Z}^\ell$. If $k_0 = 0$ then ${}^{\#}\tilde{\mathcal{P}}(M, 0) \geqslant 1$. If $k_0 \neq 0$, we extend $\breve{H}$ periodically in p to obtain a new Hamiltonian function $\hat{H}$, which satisfies Proposition 2.15. Repeating the proof of Section 4.1, we have ${}^{\#}\tilde{\mathcal{P}}(M, k_0) \geqslant n + 1$. ■

ACKNOWLEDGEMENTS. This article is a part of my Ph. D. thesis. I would like to thank my thesis advisor, Professor Yiming Long, for introducing me to this filed, drawing my attention to the work of P. Felmer, and many helpful discussions on this topic. I would also thank the referee for his/her valuable comments on the paper.

References

[1] T. Bartsch, Z. Q. Wang, Periodic Solutions of Spatially Periodic, Even Hamiltonian Systems, *J. Diff. Eqns.* 135 (1997) 103–128.

[2] T. Bartsch, Z. Q. Wang, Periodic solutions of even Hamiltonian systems on the torus, *Math. Z.* 224 (1997) 65–76.

[3] K. C. Chang, On the periodic nonlinearity and the multiplicity of solutions, *Nonlinear Analysis* 13 (1989) 527-538.

[4] C. Conley, E. Zehnder, The Birkhoff-Lewis fixed point theorem and a conjecture of V. I. Arnold, *Invent. math.* 73 (1983) 33–49.

[5] P. Felmer, Periodic Solutions of Spatially Periodic Hamiltonian Systems, *J. Diff. Eqns.* 98 (1992) 143–168.

[6] P. Felmer, Rotation Type Solutions for Spatially Periodic Hamiltonian Systems, *Nonlinear Analysis TMA* 19 (1992) 409–425.

[7] C. Golé, Monotone maps of $T^n \times \mathbb{R}^n$ and their periodic orbits, The geometry of Hamiltonian systems (ed. T. Ratin), MSRI Publications 22 (Springer, New York) (1991) 341–366.

[8] H. Hofer, E. Zehnder, *Symplectic invariants and Hamiltonian dynamics*, Birkhäuser, 1994.

[9] M. Y. Jiang, Periodic Solutions of Hamiltonian systems on hypersurfaces in a torus, *manuscripta math* 85 (1994) 307–321.

[10] F. Josellis, Lyusternik-Schnirelman Theory for Flows and Periodic Orbits for Hamiltonian systems on $T^n \times \mathbb{R}^n$, *Proc. London Math. Soc.* 68 (1994) 641–672.

[11] P. Rabinowitz, On a class of functionals invariant under a $\mathbb{Z}^n$ action, *Trans. Am. math. Soc.* 310 (1988) 303–311.

[12] A. Weinstein, On the hypotheses of Rabinowitz' periodic orbit theorems, *J. Diff. Eqns.* 33 (1979) 353–358.



Fig 1. simple pendulum



Fig 2. auxiliary function *f*



Fig 3. auxiliary function g